

Viscous relativistic hydrodynamics*



Ulrich Heinz

Department of Physics
The Ohio State University
191 West Woodruff Avenue
Columbus, OH 43210

Presented at

Extreme QCD

University of Wales, Swansea, Aug. 2-5, 2005

Based on work done in collaboration with **Asis Chaudhuri** and on the following key references:

W. Israel and J. M. Stewart, Ann. Phys. (N.Y.) **118**, 349 (1979)

A. Muronga, Phys. Rev. C **69**, 034903 (2004)

A. Muronga and D. Rischke, nucl-th/0407114

A. Chaudhuri and U. Heinz, nucl-th/0504022

*Work supported by the U.S. Department of Energy (DOE)

Starting point: The conservation laws

$$\partial_\mu N^\mu = 0 \quad \text{charge conservation}$$

$$\partial_\mu T^{\mu\nu} = 0 \quad \text{energy-momentum conservation}$$

$$\partial_\mu S^\mu \geq 0 \quad 2^{\text{nd}} \text{ law of thermodynamics}$$

Ideal fluid decomposition

Ideal fluid dynamics \Longleftrightarrow local thermal equilibrium $f(x, p) = f_{\text{eq}}(x, p)$
 \Longleftrightarrow collision time scale \ll macroscopic time scales

$$N^\mu = \int \frac{d^3p}{E} p^\mu f(x, p) = n u^\mu \quad n = (\text{net}) \text{ charge density}$$

$$T^{\mu\nu} = \int \frac{d^3p}{E} p^\mu p^\nu f(x, p) \quad e = \text{energy density}$$

$$= (e + p) u^\mu u^\nu - p g^{\mu\nu} \quad p = \text{pressure}$$

$$= e u^\mu u^\nu - p \Delta^{\mu\nu} \quad \Delta^{\mu\nu} = g^{\mu\nu} - u^\mu u^\nu$$

$$S^\mu = s u^\mu \quad s = \text{entropy density}$$

First law of thermodynamics: $Ts = p - \mu n + e$

$$\partial_\mu N^\mu = \partial_\mu T^{\mu\nu} = 0 \implies \partial_\mu S^\mu = 0$$

(in absence of shock discontinuities, entropy is conserved)

Ideal fluid equations (in comoving frame)

Convective and transverse derivative: $\partial_\mu = u^\mu D + \nabla^\mu$
 $D \equiv u^\nu \partial_\nu, \quad \nabla^\mu \equiv \Delta^{\mu\nu} \partial_\nu$

$$\dot{n} = -n \theta$$

$$\dot{e} = -(e + p) \theta$$

$$\dot{u}^\mu = \frac{\nabla^\mu p}{e + p}$$

$$p = p(n, e)$$

$\dot{f} = u^\mu \partial_\mu f \equiv Df$ = time derivative in
local rest frame

$\theta \equiv \partial \cdot u$ = local expansion rate

equation of state (EOS)

6 equations for 6 unknowns: n, e, p, u^μ

Non-ideal fluid decomposition

$$f(x, p) = f_{\text{eq}}(x, p) + \delta f(x, p)$$

$$\begin{aligned} N^\mu &= n u^\mu + V^\mu \\ &= N_{\text{eq}}^\mu + \delta N^\mu \\ T^{\mu\nu} &= e u^\mu u^\nu - p \Delta^{\mu\nu} \\ &\quad - \Pi \Delta^{\mu\nu} + \pi^{\mu\nu} \\ &\quad + W^\mu u^\nu + W^\nu u^\mu \\ &= T_{\text{eq}}^{\mu\nu} + \delta T^{\mu\nu} \\ S^\mu &= s u^\mu + \Phi^\mu \\ &= S_{\text{eq}}^\mu + \delta S^\mu \end{aligned}$$

$$n = u_\mu N^\mu$$

$$V^\mu = \Delta^{\mu\nu} N_\nu = \text{charge flow in l.r.f.}$$

$$e = u_\mu T^{\mu\nu} u_\nu$$

$$\Pi = -\frac{1}{3} \Delta_{\mu\nu} T^{\mu\nu} - p = \text{viscous bulk pressure}$$

$$W^\mu = u^\nu T_{\nu\lambda} \Delta^{\lambda\mu} = \text{energy flow in l.r.f.}$$

$$= q^\mu + \frac{e+p}{n} V^\mu \quad q^\mu = \text{heat flow in l.r.f.}$$

$$\pi^{\mu\nu} = T^{\langle\mu\nu\rangle}$$

$$\equiv \left[\frac{1}{2} (\Delta^{\mu\sigma} \Delta^{\nu\tau} + \Delta^{\mu\tau} \Delta^{\nu\sigma}) - \frac{1}{3} \Delta^{\mu\nu} \Delta^{\tau\sigma} \right] T_{\tau\sigma}$$

$$= \text{viscous shear pressure tensor } (\pi^\mu_\mu = 0)$$

$$s = u_\mu S^\mu$$

$$\Phi^\mu = \Delta^{\mu\nu} S_\nu = \text{entropy flow in l.r.f.}$$

Frame choice and matching conditions

The local equilibrium distribution $f_{\text{eq}}(x, p)$ (with local temperature $T(x)$ and chemical potential $\mu(x)$) that best matches the non-equilibrium $f(x, p)$ is defined by the **matching conditions**

$$u_\mu \delta T^{\mu\nu} u_\nu = u_\mu \delta N^\mu = 0$$

Local rest frame is ambiguous:

Eckart frame: $V^\mu = 0, \quad q^\mu = W^\mu$

Landau frame: $W^\mu = 0, \quad q^\mu = -\frac{e+p}{n} V^\mu$

(Intermediate frames also possible.)

\Rightarrow Need $1 + 3 + 5 = 9$ additional equations for $\Pi, q^\mu, \pi^{\mu\nu}$ from underlying transport theory.

Non-ideal fluid equations

$$\begin{aligned}
 \dot{n} &= -n\theta - \nabla \cdot V + V \cdot \dot{u} \\
 \dot{e} &= -(e+p+\Pi)\theta + \pi_{\mu\nu} \nabla^{\langle\mu} u^{\nu\rangle} - \nabla \cdot W + 2W \cdot \dot{u} \\
 (e+p+\Pi) \dot{u}^\mu &= \nabla^\mu(p+\Pi) - \Delta^{\mu\nu} \nabla^\sigma \pi_{\nu\sigma} + \pi^{\mu\nu} \dot{u}_\nu \\
 &\quad - [\Delta^{\mu\nu} \dot{W}_\nu + W^\mu \theta + (W \cdot \nabla) u^\mu]
 \end{aligned}$$

Depending on frame, can set either $V^\mu = 0$ or $W^\mu = 0$. In Landau frame ($W^\mu = 0$) and for baryon-free systems ($n = 0$, no heat conduction) equations simplify to:

$$\begin{aligned}
 \dot{e} &= -(e+p+\Pi)\theta + \pi_{\mu\nu} \nabla^{\langle\mu} u^{\nu\rangle} \\
 (e+p+\Pi) \dot{u}^\mu &= \nabla^\mu(p+\Pi) - \Delta^{\mu\nu} \nabla^\sigma \pi_{\nu\sigma} + \pi^{\mu\nu} \dot{u}_\nu
 \end{aligned}$$

Need extra equations for bulk and shear viscous pressures Π , $\pi^{\mu\nu}$.

Follow Chapman-Enskog strategy: write $f(x, p) = f_{\text{eq}}(p \cdot u(x); T(x), \mu(x)) + \delta f(x, p)$ and assume that $\delta f \ll f$ (and thus δN^μ and $\delta T^{\mu\nu}$) can be expanded in gradients of equilibrium parameters T, μ, u_μ .

The second law of thermodynamics (I)

In equilibrium the identity $Ts = p - \mu n + e$ can be rewritten as

$$S_{\text{eq}}^\mu = p(\alpha, \beta)\beta^\mu - \alpha N_{\text{eq}}^\mu + \beta_\nu T_{\text{eq}}^{\mu\nu}$$

where $\alpha \equiv \mu/T$, $\beta \equiv 1/T$, and $\beta^\mu \equiv u^\mu/T$.

The most general off-equilibrium generalization is (Israel & Stewart 1979)

$$S^\mu = p(\alpha, \beta)\beta^\mu - \alpha N^\mu + \beta_\nu T^{\mu\nu} + Q^\mu(\delta N^\mu, \delta T^{\mu\nu})$$

where Q^μ is second and higher order in the off-equilibrium deviations δN^μ and $\delta T^{\mu\nu}$.

The Gibbs-Duhem relation $dp = s dT + n d\mu$ can be recast as

$$\partial_\mu(p(\alpha, \beta)\beta^\mu) = N_{\text{eq}}^\mu \partial_\mu \alpha - T_{\text{eq}}^{\mu\nu} \partial_\mu \beta_\nu$$

Using also the conservation laws, the entropy production rate takes the form

$$\partial_\mu S^\mu = -\delta N^\mu \partial_\mu \alpha + \delta T^{\mu\nu} \partial_\mu \beta_\nu + \partial_\mu Q^\mu$$

The second law of thermodynamics (II)

In the **Chapman-Enskog** spirit, one now postulates linear relations between the off-equilibrium flows δN^μ , $\delta T^{\mu\nu}$ and the thermodynamic forces $\partial^\mu \alpha$, $\partial^{(\mu} \beta^{\nu)}$, consistent with the second law

$$\partial_\mu S^\mu = -\delta N^\mu \partial_\mu \alpha + \delta T^{\mu\nu} \partial_\mu \beta_\nu + \partial_\mu Q^\mu \geq 0$$

These relations depend on the choice of Q^μ . Standard dissipative relativistic fluid dynamics assumes $Q^\mu = 0$. In this case

$$T \partial_\mu S^\mu = \Pi X - q^\mu X_\mu + \pi^{\mu\nu} X_{\langle\mu\nu\rangle} \equiv \frac{\Pi^2}{\zeta} - \frac{q^\mu q_\mu}{2\lambda T} + \frac{\pi^{\alpha\beta} \pi_{\alpha\beta}}{2\eta} \geq 0,$$

with thermodynamic forces $X \equiv -\nabla \cdot u = -\theta$, $X^\mu \equiv \frac{\nabla^\mu T}{T} - \dot{u}^\mu = -\frac{nT}{e+p} \nabla^\mu \left(\frac{\mu}{T}\right)$ and $X_{\langle\mu\nu\rangle} \equiv \nabla_{\langle\mu} u_{\nu\rangle}$, can be satisfied by setting

$$\Pi = -\zeta \theta, \quad q^\mu = -\lambda \frac{nT^2}{e+p} \nabla^\mu \left(\frac{\mu}{T}\right), \quad \pi^{\mu\nu} = 2\eta \nabla^{\langle\mu} u^{\nu\rangle}$$

with **positive** transport coefficients $\zeta \geq 0$, $\lambda \geq 0$, and $\eta \geq 0$.

Unfortunately, plugging these equations for Π , q^μ , and $\pi^{\mu\nu}$ directly into the non-ideal hydro equations leads to **acausal** signal propagation.

The second law of thermodynamics (III)

Causal relativistic fluid dynamics requires keeping Q^μ in the entropy flux, at least up to terms of second order in the irreversible flows.

$$S^\mu = su^\mu + \frac{q^\mu}{T} + Q^\mu$$

Setting $q^\nu = 0$ ($n = 0$) for simplicity, we get up to second order

$$S^\mu = su^\mu - (\beta_0 \Pi^2 + \beta_2 \pi_{\nu\lambda} \pi^{\nu\lambda}) \frac{u^\mu}{2T}$$

This yields (after some algebra)

$$\begin{aligned} T \partial_\mu S^\mu &= \Pi \left[-\theta - \beta_0 \dot{\Pi} - \Pi T \partial_\mu \left(\frac{\beta_0 u^\mu}{2T} \right) \right] \\ &+ \pi^{\alpha\beta} \left[\nabla_{\langle\alpha} u_{\beta\rangle} - \beta_2 \dot{\pi}_{\alpha\beta} - \pi_{\alpha\beta} T \partial_\mu \left(\frac{\beta_2 u^\mu}{2T} \right) \right] \\ &\stackrel{!}{=} \frac{\Pi^2}{\zeta} + \frac{\pi^{\alpha\beta} \pi_{\alpha\beta}}{2\eta} \geq 0 \end{aligned}$$

The thermodynamic forces $-\theta$, $\nabla_{\langle\alpha} u_{\beta\rangle}$ are seen to be self-consistently modified by the irreversible flows Π , $\pi_{\alpha\beta}$. This leads to dynamical (“transport”) equations for Π , $\pi_{\alpha\beta}$.

Transport equations for the irreversible flows

The resulting transport equations for Π , $\pi_{\alpha\beta}$ are

$$\begin{aligned}\dot{\Pi} &= -\frac{1}{\tau_{\Pi}} \left[\Pi + \zeta\theta + \Pi\zeta T\partial_{\mu} \left(\frac{\tau_{\Pi}u^{\mu}}{2\zeta T} \right) \right] \approx -\frac{1}{\tau_{\Pi}} [\Pi + \zeta\theta] \\ \dot{\pi}_{\alpha\beta} &= -\frac{1}{\tau_{\pi}} \left[\pi_{\alpha\beta} - 2\zeta\nabla_{\langle\alpha}u_{\beta\rangle} + \pi_{\alpha\beta}\eta T\partial_{\mu} \left(\frac{\tau_{\pi}u^{\mu}}{2\eta T} \right) \right] \\ &\approx -\frac{1}{\tau_{\pi}} [\pi_{\alpha\beta} - 2\zeta\nabla_{\langle\alpha}u_{\beta\rangle}]\end{aligned}$$

where we introduced the **relaxation times** $\tau_{\Pi} = \zeta\beta_0$, $\tau_{\pi} = 2\eta\beta_2$. In principle, both ζ , η and τ_{Π} , τ_{π} should be calculated from the underlying kinetic theory.

The **purple terms** are of second order in the derivatives of the thermodynamic equilibrium quantities and, in the regime of validity of the approach, should be neglected relative to the other terms.

[Keeping them would require also keeping third-order terms in the entropy flow Q^{μ} for consistency, and would modify both the effective local relaxation times $\tau_{\Pi,\pi}$ and the viscosities η , ζ by amounts which depend on the local hydrodynamic expansion rate.]

Viscous relativistic hydrodynamics (Israel & Stewart 1979)

Include shear viscosity η , neglect bulk viscosity (massless partons) and heat conduction ($\mu_B \approx 0$); solve

$$\partial_\mu T^{\mu\nu} = 0$$

with modified energy momentum tensor

$$T^{\mu\nu}(x) = (e(x) + p(x))u^\mu(x)u^\nu(x) - g^{\mu\nu}p(x) + \pi^{\mu\nu}.$$

$\pi^{\mu\nu}$ = traceless viscous pressure tensor which relaxes locally to 2η times the shear tensor $\nabla^{\langle\mu} u^{\nu\rangle}$ on a microscopic kinetic time scale τ_π :

$$D\pi^{\mu\nu} = -\frac{1}{\tau_\pi} (\pi^{\mu\nu} - 2\eta \nabla^{\langle\mu} u^{\nu\rangle})$$

where $D \equiv u^\mu \partial_\mu$ is the time derivative in the local rest frame.

Kinetic theory relates η and τ_π , but for a strongly coupled QGP neither η nor this relation are known \implies treat η and τ_π as independent phenomenological parameters. For consistency: $\tau_\pi \theta \ll 1$ ($\theta = \partial^\mu u_\mu$ = local expansion rate).

(1+1)-d viscous hydrodynamic equations

(Muronga & Rischke 2004, Chaudhuri & Heinz 2005)

Azimuthally symmetric transverse dynamics with long. boost invariance:
Use (τ, r, ϕ, η) coordinates and solve

- hydrodynamic equations for $T^{\tau\tau} = (e + \mathcal{P})\gamma_r^2 - \mathcal{P}$, $T^{\tau r} = (e + \mathcal{P})\gamma_r^2 v_r$
(with “effective pressure” $\mathcal{P} = p - r^2 \pi^{\phi\phi} - \tau^2 \pi^{\eta\eta}$) together with
- kinetic relaxation equations for $\pi^{\phi\phi}$, $\pi^{\eta\eta}$:



$$\begin{aligned}\frac{1}{\tau} \partial_\tau (\tau T^{\tau\tau}) + \frac{1}{r} \partial_r (r (T^{\tau\tau} + \mathcal{P}) v_r) &= - \frac{p + \tau^2 \pi^{\eta\eta}}{\tau}, \\ \frac{1}{\tau} \partial_\tau (\tau T^{\tau r}) + \frac{1}{r} \partial_r (r (T^{\tau r} v_r + \mathcal{P})) &= + \frac{p + r^2 \pi^{\phi\phi}}{r}, \\ (\partial_\tau + v_r \partial_r) \pi^{\eta\eta} &= - \frac{1}{\gamma_r \tau_\pi} \left[\pi^{\eta\eta} - \frac{2\eta}{\tau^2} \left(\frac{\theta}{3} - \frac{\gamma_r}{\tau} \right) \right], \\ (\partial_\tau + v_r \partial_r) \pi^{\phi\phi} &= - \frac{1}{\gamma_r \tau_\pi} \left[\pi^{\phi\phi} - \frac{2\eta}{r^2} \left(\frac{\theta}{3} - \frac{\gamma_r v_r}{r} \right) \right].\end{aligned}$$

Close equations with EOS $p(e)$ where $e = T^{\tau\tau} - v_r T^{\tau r}$ and $v_r = T^{\tau r} / (T^{\tau\tau} + \mathcal{P})$.

(1+1)-d viscous hydrodynamics: first results (I)

(Chaudhuri & Heinz, nucl-th/0504022)

Viscosity effects on freeze-out surface ($\tau_\pi = \frac{3\eta}{2p}$, $\pi_{ini}^{rr} = \frac{2\eta}{3\tau_i}$):



- Both sets of initial conditions have similar initial total entropy.
- Viscosity slows down cooling and gives more time for transverse expansion.
- Viscous effects are larger for smaller τ_i , due to faster initial expansion rate.

(1+1)-d viscous hydrodynamics: first results (II)




(Chaudhuri & Heinz, nucl-th/0504022)

Sensitivity to initial π^{rr} , $\frac{\eta}{s}$, and relaxation time τ_π ($T_f = 160 \text{ MeV}$):

$$\tau_\pi = \frac{3\eta}{2p}, \quad \frac{\eta}{s} = 0.135$$

$$\tau_\pi = \frac{3\eta}{2p}, \quad \pi_{\text{ini}}^{rr} = \frac{2\eta}{3\tau_i}$$

$$\eta/s = 0.135, \quad \pi_{\text{ini}}^{rr} = \frac{2\eta}{3\tau_i}$$



- Larger initial viscous pressures create larger overall viscous effects (“memory effect”)
- Significant viscous effects for $\frac{\eta}{s} > \frac{\hbar}{4\pi}$
- At fixed $\frac{\eta}{s}$, viscous effects increase with increasing relaxation time τ_π

Summary

- Causal relativistic dissipative hydrodynamics requires solution of a coupled set of (i) **hydrodynamic equations** with additional irreversible flow corrections and (ii) **kinetic relaxation equations** for these irreversible flows.
- Relaxation equations for dissipative flows are derived from a **second-order approach** to implementing the 2nd law of thermodynamics, keeping terms up to second order in derivatives of equilibrium quantities.
- For **each** non-vanishing space-time component of the hydro equations, we must solve **one** transport equation for **each** non-vanishing transport coefficient (bulk viscosity, shear viscosity, heat conduction) \Rightarrow **significantly increased numerical complexity**.
- (1+1)-dimensional viscous hydro is under investigation;
(2+1)-dimensional code is under construction.